

# Intrahepatic Glissonian Approach for Laparoscopic Right Trisectionectomy

Marcel A.C. Machado, MD, Fabio F. Makdissi, Rodrigo C.T. Surjan,  
Andre C. Oliveira, Victor F. Pilla, and A.R. Teixeira

## Introduction

**L**IVER RESECTION is the definitive treatment for several benign and malignant liver diseases. Experience with laparoscopic procedures and recent advances in laparoscopic devices have created an evolving interest in the application of these techniques to liver resection.<sup>1-3</sup> However, laparoscopic liver resections may be technically demanding. Pedicle control is an important step of liver resection. Anatomic hepatectomies usually require extensive hilar dissection. To facilitate pedicle control and reduce operating time, we have previously described a technique for laparoscopic right-liver resections by using the intrahepatic Glissonian approach.<sup>4</sup> Laparoscopic right-liver trisectionectomy is a very complex procedure, and, to our knowledge, there is only one technical description, so far, in the English literature.<sup>5</sup> This video demonstrates technical aspects of a totally laparoscopic right trisectionectomy, using the intrahepatic Glissonian approach. Our technique differs from that previously described<sup>5</sup> by intrahepatic pedicle control and total control of venous outflow, which makes the procedure easier and safer for laparoscopy.

## Patient and Method

A 22-year-old woman with a giant angiomyolipoma was referred for surgical treatment. The patient was placed in a left semilateral decubitus position with the surgeon between the patient's legs. Five trocars (three 12 and two 5 mm) were used. The pneumoperitoneum is established at a pressure of 12 mm Hg. Round and falciform ligaments are taken down close to the abdominal wall in order to facilitate left-liver fixation at the end of the procedure. The falciform and coronary ligaments are divided by using laparoscopic coagulation shears (LCS; Ethicon Endo-Surgery Industries, Cincinnati, OH) to expose the suprahepatic inferior vena cava. After cholecystectomy, the right hepatic artery is ligated, resulting in an ischemic delineation of the right liver. Due to previous right-portal-vein embolization in this patient, the hepatic pedicle was not fully dissected. The right liver is then fully mobilized, and the inferior vena cava is dissected. A large inferior right hepatic vein arising from segment 6 is ligated and divided

between metallic clips. Another accessory right hepatic vein from segment 7 (middle-right hepatic vein) is divided with a vascular endoscopic stapler. The right hepatic vein is finally encircled, and downward retraction permits the safe application of a vascular endoscopic stapler. The stapler is fired, leaving three lines of metallic clips. With this maneuver, the anterior surface of the retrohepatic vena cava is completely exposed. The main trunk, including the middle and left hepatic veins, is now the only venous drainage of the liver. It is encircled and traction or temporary clamping permits complete outflow control of the liver, minimizing bleeding during liver transection. At this time, the intrahepatic access to the main right Glissonian pedicle is achieved with two small incisions: An incision is performed on the right portion of the caudate lobe and another anterior incision is made in front of the hilum. An endoscopic vascular-stapling device is inserted between these incisions, and the stapler is fired. All these steps are performed without the Pringle maneuver and without hand assistance.

The line of liver transection is marked along the liver surface, including segment 4. To avoid possible damage to pedicles from segments 2 and 3, the line of transection should be placed 1 cm right from the falciform ligament. The division of the liver parenchyma should be performed under central venous pressure as low as possible. The Glissonian pedicle from segment 4 is divided during liver transection. Liver transection should be performed toward the main trunk to prevent damage to the left hepatic vein. Liver transection is accomplished with a Harmonic Scalpel and endoscopic stapling device, as appropriate. The specimen is extracted through a suprapubic incision, and the pneumoperitoneum is reestablished. Raw surface area is then checked for hemostasia and biliary leakage, and absorbable hemostat tissue (Surgicel; Ethicon Industries, Cincinnati, OH) is applied. The falciform ligament is then fixed to the abdominal wall in order to prevent the remnant liver from rotating spontaneously into the right subphrenic space and causing left-hepatic-vein kinking.<sup>6</sup> One round 19-F Blake abdominal drain (Ethicon) is left in place. The right hepatic trisectionectomy is then completed.

## Results

Operative time was 360 minutes, and hospital stay was 7 days. Apart from self-limited biliary leakage, postoperative recovery was uneventful.

## Conclusions

Totally laparoscopic right trisectionectomy is safe and feasible in selected patients and should be considered for patients with benign or malignant liver neoplasms. The described technique, with the use of the intrahepatic Glissonian approach and control of venous outflow, may facilitate laparoscopic extended liver resections by reducing the technical difficulties in pedicle control and may diminish bleeding during liver transection.

## Disclosure Statement

No competing financial interests exist.

## References

1. Soubrane O, Cherqui D, Scatton O, Stenard F, Bernard D, Branchereau S, Martelli H, Gauthier F. Laparoscopic left lateral sectionectomy in living donors: Safety and reproducibility of the technique in a single center. *Ann Surg* 2006; 244:815–820.
2. O'Rourke N, Fielding G. Laparoscopic right hepatectomy: Surgical technique. *J Gastrointest Surg* 2004;8:213–216.
3. Koffron AJ, Auffenberg G, Kung R, Abecassis M. Evaluation of 300 minimally invasive liver resections at a single institution: Less is more. *Ann Surg* 2007;246:385–392.
4. Machado MA, Makdissi FF, Galvão FH, Machado MC. Intrahepatic Glissonian approach for laparoscopic right segmental liver resections. *Am J Surg* 2008;196:e38–e42.
5. Gumbs AA, Gayet B. Totally laparoscopic extended right hepatectomy. *Surg Endosc* 2008;22:2076–2077.
6. Ogata S, Kianmanesh R, Belghiti J. Doppler assessment after right hepatectomy confirms the need to fix the remnant left liver in the anatomical position. *Br J Surg* 2005;92:592–595.

Address correspondence to:

*Marcel A.C. Machado, MD*

*LIM-37*

*Department of Gastroenterology*

*University of São Paulo*

*Rua Evangelista Rodrigues 407*

*São Paulo 05463-000*

*Brazil*

*E-mail: dr@drmarcel.com.br*